

Comparison of two mesoscale LAM-EPS generation methods for the prediction of heavy rains over the Western Mediterranean: the HyMeX IOP8 event.

Maria del Mar Vich,

Octavio Jaume, Víctor Homar and Romualdo Romero

Meteorology Group, Universitat de les Illes Balears, Spain



Why the IOP8 event?

28-30 September 2012

- **Heavy precipitation:**
 - 28 - Andalusia and Murcia
 - 29 - Catalonia, Valencia and Balearic Islands
 - 30 - Golf of Lyon
- **Personal losses:** 11 deaths
- **Material losses:** 120M€



Why the IOP8 event?

28-30 September 2012

- **Heavy precipitation:**
 - 28 - Andalusia and Murcia
 - 29 - Catalonia, Valencia and Balearic Islands
 - 30 - Golf of Lyon
- **Personal losses:** 11 deaths
- **Material losses:** 120M€



Why the IOP8 event?




Why the IOP8 event?



Why the IOP8 event?

ARPEGE analysis


27 September 2012 at 00 UTC



Why the IOP8 event?

ARPEGE analysis


28 September 2012 at 00 UTC



Why the IOP8 event?

ARPEGE analysis


29 September 2012 at 00 UTC



Why the IOP8 event?

ARPEGE analysis

30 September 2012 at 00 UTC




Why the IOP8 event?

24h Accumulated rainfall from AEMET stations


28 September 2012

29 September 2012


RR OBSERVATION 2012/9/28 [00UTC(J) to 00UTC(J+1)]
(mm/24hr)



RR OBSERVATION 2012/9/29 [00UTC(J) to 00UTC(J+1)]
(mm/24hr)



Which two mesoscale LAM-EPSs?



Which two mesoscale LAM-EPSs?

The **PV-gradient** ensemble

To perturb the initial and boundary conditions
by
perturbing the 3-D structure of the PV field

- Why to perturb the **PV** field?
 - ① PV inversion technique → perturb the T and Wind fields
 - ② precursor upper-level PV structures → mid-latitude cyclonic situations
- Perturb: *how much* and *where*?

How much?

PV error climatology

Comparing the PV fields of

ECMWF **analysis** $\longleftrightarrow$ ECMWF **24 h forecast**,


of a large collection of **MEDEX cyclones**, one can define:

- The **displacement error** (DE): the minimum displacement of the 24 h forecast PV field showing local maximum correlation with the analysis PV field
- The **intensity error** (IE): the difference between the displaced 24 h forecast PV field and analysis PV field relative to the analysis PV average


MEDEX: Mediterranean Experiment on Cyclones that produce High Impact Weather in the Mediterranean

Where?

The most intense values and gradients PV zones at 300 hPa



MM5



Initial and Boundaries
conditions

“by”

ECMWF analyses

Which two mesoscale LAM-EPSs?

The ECWF global EPS


- Perturbs **initial and boundary conditions** with a combination of **singular vectors**, computed to optimize total energy growth over a 48h time interval.
- Runs with a horizontal resolution of **32 km** during the first 10 days of the EPS and of 63 km during the extension beyond day 10.
- Has 50 perturbed members plus the non-perturbed one. Total: **51 members**.

Which two mesoscale LAM-EPSs? ECMWF EPS downscaling



Dynamical downscaling of the ECWF global EPS

“by”

WRF



Which two mesoscale LAM-EPSs?



Numerical Model	MM5	WRF
Number of members	51	51
IC generation method	Perturb over the most intense values and gradients PV zones	Perturbing with a combination of singular vectors, computed to optimize total energy growth over a 48h time interval

Already tested
BASELINE

And the “winner” is?



ECMWF
downscaling

PV-gradient

And the “winner” is?

I will let you know!!

